



# Historic Trends & Future Climatic Projections for VT



---

**Dr. Alan K. Betts**

**Atmospheric Research, Pittsford, VT 05763**

**[akbetts@aol.com](mailto:akbetts@aol.com)**  
**<http://alanbetts.com>**


---

***VULNERABILITY ASSESSMENT WORKSHOP***  
***Montpelier, VT***

**July 9, 2012**


# Climate of Vermont

- Climate is a mean (10-30y)
- $T_{\max}$ ,  $T$ ,  $T_{\min}$
- Large seasonal range
- *Freezing T of water critical to climate*



# Diurnal Temperature Range

- $T_{\max} - T_{\min}$
- Mean daily range of T **varies with season**
- Related to RH and  $LW_{\text{net}}$





- Earth sustains life
- Weather changes fast
- Climate changes slowly
- Greenhouse gases keep Earth warm
- Burning fossil fuels – coal, oil and gas – is having a big effect on climate by increasing greenhouse gases: CO<sub>2</sub> and H<sub>2</sub>O




***January 2, 2012: NASA***

# What Is Happening to Vermont?

- **PAST 40/50 years** (anthropogenic forcing detectible)
- **Warming twice as fast in winter than summer**
- **Winter severity decreasing**
- **Lakes frozen less by 6.9 ( $\pm 1.5$ ) days / decade**
- **Growing season longer by 3.7 ( $\pm 1.1$ ) days / decade**
- **Spring coming earlier by 2-3 days / decade**
- **Extremes increasing**
- *Evaporation increases with  $T$*
- *More 'stationary weather patterns'*


# Vermont Temperature Trends 1961-2008

- **Summer  $+0.4^{\circ}\text{F}$  / decade**
- **Winter  $+0.9^{\circ}\text{F}$  / decade**
- **Larger variability, larger trend**
- ***Less snow (and increased water vapor) drive larger winter warming***



*Note: trends since 1961: early 1950's warmer. Trends for last 4-5 decades consistent with model projections for the next few decades*


# Slope Depends on Start-Date



**Stager and Thill (2010) chose 1975. I have gone back to 1961**



# Attribution for temperature change (AR4, Chapter 9, p.703)




- **Blue: natural forcings**
- **Red: natural & anthropogenic forcings**
- **Separate around 1960-1970** (*Solid line is data: Bands represent 90% of models*)



# Global Temperature Rise 1880 – Present



2100: +5°F



NASA-GISS, 2011

# Lake Freeze-up & Ice-out Changing


## Frozen Period Shrinking Fast




- Ice-out earlier by 2.9 ( $\pm 1.0$ ) days / decade
- Freeze-up later by 3.9 ( $\pm 1.1$ ) days / decade
- Rivers and soils similar?

# Winter Hardiness Zones - Northeast

# Change in 16 years



## 1990



2006

### Minimum winter T

#### 4: -30 to -20°F

## 5: -20 to -10°F

6: -10 to 0°F

Zone




## USDA Hardiness Zones

© 2006 by The National Arbor Day Foundation®




# Latest detailed map

- **USDA : VT Hardiness Zone Map 1976-2005 [mean 1990]**
- **A trend of half a zone in 16-20 years is +2.5-3.1°F/decade [triple rise of winter mean]**
- <http://planthardiness.ars.usda.gov/PHZMWeb/>





# Heating Degree Days and Days below 0°F (Burlington)

- HDD trend  
289 ( $\pm 37$ )  
/decade
- $T_{\min} < 0^{\circ}\text{F}$   
4.1 ( $\pm 0.7$ )  
days  
/decade







# Lilac Leaf and Bloom




- Leaf-out -2.9 days/decade; Bloom -1.6 days/decade
- Large year-to-year variation related to temperature: 4 to 5 days/ °C

# Lilac Leaf-out and Ice-out Coupled



- Lilac leaf and lake ice-out both depend on Feb. Mar. and April temperatures
- Trends indicate earlier spring

# First and Last Frosts Changing



- Growing season for frost-sensitive plants increasing **3.7 ( $\pm 1.1$ ) days / decade**
- A help for growing “local food”



# Shrinking Winter: Pittsford, VT

(Freeze-up used to be mid-November)



**January 7, 2007**

**December 2006:**

- Warmest on record



**January 10, 2008**

**Warm Fall:**

- Record Arctic sea-ice melt
- Snow cover in December, ground unfrozen



# January 2, 2012




# March 11, 2012



## October 2011– March 2012

- **Warmest 6 months on record**
- **My garden frozen only 67 days**
- **No permanent snow cover west of Green Mntns**
- **Contrast snowy winter 2010-11**

Oct 2011-Mar 2012 Statewide Ranks  
National Climatic Data Center/NESDIS/NOAA





# Early Spring: Daffodils, Forsythia

*79°F on March 22, 2012*



*Pittsford Vermont*

*3/22/12*



*Pittsford Vermont*


*3/24/12*

# Temperature and Precipitation Maps

<http://www.ncdc.noaa.gov/temp-and-precip/maps.php>


## March-May 2012 Statewide Ranks

National Climatic Data Center/NESDIS/NOAA



## March-May 2012 Statewide Ranks

National Climatic Data Center/NESDIS/NOAA



- Visual grasp of spring 2012 climate extreme
- Continental-scale patterns: Records keep falling
- Note fires in dry Colorado




# Vermont Winter 2006



- Sun is low; snow reflects sunlight, except where there are trees - shadows
- Sunlight reflected, stays cold; little evaporation, clear sky; earth cools to space
- ***Positive feedback: Less snow, warmer winters (2012)***

# Arctic Sea Ice Loss Has Accelerated



- **Positive feedbacks speed melting**
- **Less ice, less sunlight reflected**
- **More evaporation, larger water vapor greenhouse effect**

- **Record ice loss in 2007** ([www.nsidc.org](http://www.nsidc.org))




- most ice now thin and only 1-2 years old

- **Open water in Oct. Nov. favors warmer Fall**

*At the end of Nov. Hudson Bay was still nearly ice-free.*

# Sea Ice Trends


- Sea ice is **thinning rapidly**
- Observed September decline appears to be **faster than IPCC-AR4 climate model projections**
- *AR5 projections should be faster!*






# June snow cover minimum

Northern Hemisphere Snow Cover Anomaly  
June 2012



Northern Hemisphere Snow Cover Anomaly  
June 1967 - 2012



- New minimum by  $10^6$  km<sup>2</sup> (1971-2000 ref)

# Spring Climate Transition



- **Before leaf-out**

**Little evaporation** → Dry atmosphere, low humidity  
→ Low water vapor greenhouse  
→ Large cooling at night  
→ Large diurnal temp. range  
giving warm days, cool nights and frost

- **After leaf-out**

**Large evaporation** → Wet atmosphere, low cloudbase  
→ Small cooling at night  
→ Reduced maximum temperature  
→ Reduced chance of frost

- ***Spring is coming earlier***

# Summer dry-down

- Wet in spring
- Soil moisture falls: summer dry-down
- Low humidity & little rain
- *Can lock-in drought in central US*



# Recently Many Wet Summers in Vermont




- 2004, 2006, 2008, 2009, (2010), 2011 all wet
- **Direct fast evaporation off wet canopies**
- *Positive evaporation-precipitation feedback, coupled to synoptic system frequency*



# Cooling Degree Days and Days $T_{\max} > 90^{\circ}\text{F}$

- CDD trend 37 ( $\pm 11$ ) /decade
- $T_{\max} > 90^{\circ}\text{F}$  no trend - 2004-2011 were wet
- 2002 last summer with 'average' rain





# Fall Climate Transition


- **Vegetation postpones first killing frost**
- Deciduous trees still evaporating: moist air with clouds
- Water vapor & cloud greenhouse reduces cooling at night and prevents frost
- Till one night, dry air advection from north gives first hard frost.
- Vegetation shuts down, leaves turn, skies become clearer and frosts become frequent
- *The opposite of what happens in Spring with leaf-out!*



*Clear dry blue sky after frost. Forest evaporation has ended; water vapor greenhouse is reduced, so Earth cools fast to space at night*


***Later frost: Growing season getting longer***

# Carbon Dioxide Is Increasing



# 2009 Was “Good” for the Earth

## Fossil Fuel Emissions: Actual vs. IPCC Scenarios





# Rise of Greenhouse Gases (GHG) Shift Energy Balance of Planet

- The atmosphere is **transparent to light** from the sun, but **not to infrared radiation** from the earth
- **GHG:**  $\text{H}_2\text{O}$ ,  $\text{CO}_2$ ,  $\text{CH}_4$ ,  $\text{O}_3$ , CFCs absorb and reradiate IR from the surface, giving climate suitable for life by warming planet  $30^\circ\text{C}$
- $\text{CO}_2$  rise alone has a small warming effect

**BUT...**



# Water Phase Changes Give Positive Radiative Feedbacks

- As Earth warms, evaporation and water vapor increase and this is 3X amplifier on CO<sub>2</sub> rise
- As Earth warms, snow and ice decrease and reduced SW reflection amplifies warming in Arctic in summer and mid-latitudes in winter
- Doubling CO<sub>2</sub> will warm globe about 3°C (5°F)
  - Much more in the North and over land, which responds faster than oceans
  - *(also transpiration coupling of vapor and CO<sub>2</sub> fluxes)*

# Climate Change Projections


- **IPCC 2007 (Fourth Assessment - AR4)**
- *AR5 in progress with improved models – expect no large change*
- **Higher resolution: Improved aerosols, sea-ice & carbon cycle (probably slightly increased climate sensitivity and wider range between models)**



# Predicted Change in Temperature


2020-2029 and 2090-2099, relative to 1980-1999 (°C)

**“Committed”**



**(We did  
nothing for  
the last 20  
years)**

**Still up to us!**



**(We could  
halve this if  
we act now)**




**[°C]**





# Climate Model Predictions

Temperature A1B: 2080-2099




DJF Precipitation A1B: 2080-2099

DJF





Temperature A1B: 2080-2099



JJA Precipitation A1B: 2080-2099


JJA





# USGCRP (2009) - p29

- Mean of 16 CMIP3 models




The maps on this page and the previous page are based on projections of future temperature by 16 of the Coupled Model Intercomparison Project Three (CMIP3) climate models using two emissions scenarios from the Intergovernmental Panel on Climate Change (IPCC), *Special Report on Emission Scenarios* (SRES).<sup>91</sup> The "lower" scenario here is B1, while the "higher" is A2.<sup>91</sup> The brackets on the thermometers represent the likely range of model projections, though lower or higher outcomes are possible. Additional information on these scenarios is on pages 22 and 23 in the previous section, *Global Climate Change*. These maps, and others in this report, show projections at national, regional, and sub-regional scales, using well-established techniques.<sup>110</sup>



# USGCRP (2009) – p31


- More confidence where shaded
- Northern areas wetter



The maps show projected future changes in precipitation relative to the recent past as simulated by 15 climate models. The simulations are for late this century, under a higher emissions scenario.<sup>21</sup> For example, in the spring, climate models agree that northern areas are likely to get wetter, and southern areas drier. There is less confidence in exactly where the transition between wetter and drier areas will occur. Confidence in the projected changes is highest in the hatched areas.

# Observed Changes: Total Precip and Heavy Precip. (upper 1%)

Observed Change in Annual Average Precipitation  
1958 to 2008




Percent Change

<-40 -35 -30 -25 -20 -15 -10 -5 0 5 10 15 20 25 30 35 >40

While U.S. annual average precipitation has increased about 5 percent over the past 50 years, there have been important regional differences as shown above.

Increases in Amounts of Very Heavy  
Precipitation (1958 to 2007)



Updated from Groisman et al.<sup>113</sup>

The map shows percent increases in the amount falling in very heavy precipitation events (defined as the heaviest 1 percent of all daily events) from 1958 to 2007 for each region. There are clear trends toward more very heavy precipitation for the nation as a whole, and particularly in the Northeast and Midwest.

# USGCRP Northeast.pdf (2009)

- Since 1970, the annual average temperature in the Northeast has increased by 2°F, with winter temperatures rising twice this much
- Warming has resulted in many other climate-related changes including:
  - More frequent days with temperatures above 90°F
  - A longer growing season
  - Increased heavy precipitation
  - Less winter precipitation falling as snow and more as rain
  - Reduced snowpack
  - Earlier breakup of winter ice on lakes and rivers
  - Earlier spring snowmelt resulting in earlier peak river flows
  - Rising sea surface temperatures and sea level

*<http://www.globalchange.gov/publications/reports/scientific-assessments/us-impacts/regional-climate-change-impacts/northeast>*




# USGCRP Northeast.pdf (2009)

- **Over the next several decades**, temperatures in the Northeast are projected to rise an additional 2.5 to 4°F in winter and 1.5 to 3.5°F in summer.
- By mid-century and beyond, however, today's emissions choices would generate starkly different climate futures; the lower the emissions, the smaller the climatic changes and resulting impacts.
- **By late this century, under a higher emissions scenario:**
  - **Winters in the Northeast are projected to be much shorter with fewer cold days and more precipitation.**
  - **The length of the winter snow season would be cut in half across northern New York, Vermont, New Hampshire, and Maine, and reduced to a week or two in southern parts of the region.**
  - **Cities** that today experience few days above 100°F each summer would average 20 such days per summer, while certain cities, such as Hartford and Philadelphia, would average nearly 30 days over 100°F.
  - Short-term (one- to three-month) droughts are projected to occur as frequently as once each summer in the Catskill and Adirondack Mountains, and across the New England states.
  - **Hot summer conditions would arrive three weeks earlier and last three weeks longer into the fall.**
  - Sea level in this region is projected to rise more than the global average

# Vermont's Future with High and Low GHG Emissions

What  
about  
skiing?

What  
about  
tropics?



NECIA,  
2007

# Sea-level Rise Will Eventually Flood Coastal Cities

- Late 20<sup>th</sup>-century sea-level rise: 1 foot / century
- 21<sup>st</sup> century: Likely to triple to 3 - 4 feet / century
  - And continue for centuries (accelerating for business as usual)
- *<http://www.nature.com/news/us-northeast-coast-is-hotspot-for-rising-sea-levels-1.10880>*

# Many Challenges Face Us

- **Extreme weather: Floods, fires, & drought**
  - **32 weather disasters >\$1B in 2011**
- **Melting Arctic and permafrost—methane release is positive feedback**
- **Ecosystem collapse, including perhaps forest and ocean ecosystems**
- **Collapse of unsustainable human population**



# Extreme Weather (precip.)


- *Precip. is condensation of atmospheric water vapor (large latent heat release)*
- *Saturation vapor pressure at cloud-base increases steeply with temperature (6%/°C)*
- *More latent heat organizes storms, increasing convergence of vapor*
- *Quasi-stationary large-scale flow means longer rain events in low-pressure convergent regions, and longer droughts in high-pressure divergent regions*
- *As climate changes, quasi-stationary large-scale modes appear to be more frequent*
- *Wet surface: more evaporation and runoff*

# 2011 Vermont Floods

- Record spring flood on Lake Champlain
- Record floods following TS Irene
- Record wet March-August, 2011: OH to VT  
(but record drought in TX & NM)
- *Quasi-stationary pattern for 6 mos*


## March-August 2011 Statewide Ranks

National Climatic Data Center/NESDIS/NOAA



## March-August 2011 Statewide Ranks

National Climatic Data Center/NESDIS/NOAA



# Winooski River 2011

- **Two classic VT flood situations**
- **Spring flood: heavy rain and warm weather, melting large snowpack**
  - 70F (4/11) and 80F(5/27) + heavy rain
  - record April, May rainfall: **3X at BTV**
- **Irene flood: tropical storm** moved up east of Green Mountains dumping 6ins rain on wet soils (Floyd on 9/17/1999 had similar rain - but with dry soils there was less flooding)

# Discussion

- This talk <http://alanbetts.com/research>
- VTCCAdaptClimateChangeVTBetts10-29.pdf  
<http://www.anr.state.vt.us/anr/climatechange/Adaptation.html>
- *Vermont Climate Change Indicators*
- *Seasonal Climate Transitions in New England*



## Changes on NH Climate expected by 2100.

Assumptions to be used when assessing habitat and species vulnerability.

Stressor	Indicator	Changes expected by 2100 <sup>1</sup>
Direct temperature effects	Temperature increase in winter	Low: Increases 5-8°F. High: Increases 9-13°F.
	Temperature increase in summer	Low: Increases 3-7°F. High: Increases 6-14°F.
	# of days above 90° (10 now)	Low: 30 days. High: 70 days.
	# of days above 100°	Low: 6 days. High: 20 days.
	Days with snow	Low: Decreases 33%. High: Decreases 50%.
	River ice-out	Earlier by 11-13 days
Changes in precipitation	Winter precipitation	Increases 20-30% with a higher percentage as rain.
	Frequency of heavy rains	Increased
	Summer drought	Increased frequency of 1-3 month droughts, becoming annual under high emissions scenario.
Hydrology	Change in stream flow	More headwater streams become intermittent during summer months. Reduced summer flow in most rivers
Fire	Change in fire frequency	Higher temperatures and more drought events lead to increased fire frequency
Wind	Change in wind intensity	More frequent and more intense storms lead to higher frequency of damaging wind events.
Sea level rise	Sea level rise by expansion	Low: 31". High: 75".
Ocean acidification/salinity change	Changes in pH/salinity of seawater	Decrease in pH. Salinity varies depending on freshwater runoff and latitude.
Human response to climate change	Infrastructure changes	Seawalls, dams, culverts, wind power, transmission lines and other changes in developed and undeveloped landscapes change habitat permeability and alter habitat type and quality.
Altered species interactions	Growing season length	Up to 43 days longer
	First leaf	Earlier by 6.7-15 days
	Lilac bloom	Earlier by 6.3-16 days

- Highlights: ice-out and first leaf: probable 14-36 days earlier
- Sea-level misprint: was cm for expansion (?); now USGS 3-4ft with ice-melt